

CBSE Class-12 Subject Chemistry

NCERT Solutions

Chapter – 10

Haloalkanes and Haloarenes

In-text Question

1. Write structures of the following compounds:

(i) 2-Chloro-3-methylpentane


(ii) 1-Chloro-4-ethylcyclohexane

(iii) 4-tert. Butyl-3-iodoheptane

(iv) 1,4-Dibromobut-2-ene


(v) 1-Bromo-4-sec. butyl-2-methylbenzene

Ans. (i)




2-Chloro-3-methyl pentane

(ii)




1-Chloro-4-ethylcyclohexane

(iii)




4-tert-Butyl-3-iodoheptane

(iv)



1, 4-Dibromobut-2-ene

(v)




1-Bromo-4-sec-butyl-2-methylbenzene

2. Why is sulphuric acid not used during the reaction of alcohols with KI?

Ans. In the presence of sulphuric acid (H_2SO_4), KI produces HI



Since H_2SO_4 is an oxidizing agent, it oxidizes HI (produced in the reaction to I_2).




As a result, the reaction between alcohol and HI to produce alkyl iodide cannot occur.

Therefore, sulphuric acid is not used during the reaction of alcohols with KI. Instead, a non-oxidizing acid such as H_3PO_4 is used.

3. Write structures of different dihalogen derivatives of propane.


Ans. There are four different dihalogen derivatives of propane. The structures of these derivatives are shown below.

(i)




1, 1-Dibromopropane

(ii)



2, 2-Dibromopropane

(iii)



1, 2-Dibromopropane

(iv)



1, 3-Dibromopropane

4. Among the isomeric alkanes of molecular formula C_5H_{12} , identify the one that on


photochemical chlorination yields

(i) A single monochloride.

(ii) Three isomeric monochlorides.

(iii) Four isomeric monochlorides.


Ans. (i) To have a single monochloride, there should be only one type of H-atom in the isomer of the alkane of the molecular formula C_5H_{12} . This is because, replacement of any H-atom leads to the formation of the same product. The isomer is neopentane.




Neopentane

(ii) To have three isomeric monochlorides, the isomer of the alkane of the molecular formula C_5H_{12} should contain three different types of H-atoms.

Therefore, the isomer is *n*-pentane. It can be observed that there are three types of H atoms labelled as *a*, *b* and *c* in *n*-pentane.




(iii) To have four isomeric monochlorides, the isomer of the alkane of the molecular formula C_5H_{12} should contain four different types of H-atoms. Therefore, the isomer is 2-methylbutane. It can be observed that there are four types of H-atoms labelled as *a*, *b*, *c*, and *d* in 2-methylbutane.




5. Draw the structures of major monohalo products in each of the following reactions:


(i)




(ii)



(iii)




(iv)




(v)




(vi)




Ans. (i)




(ii)



(iii)




(iv)



(v)



(vi)




6. Arrange each set of compounds in order of increasing boiling points.

(i) Bromomethane, Bromoform, Chloromethane, Dibromomethane.

(ii) 1-Chloropropane, Isopropyl chloride, 1-Chlorobutane.

Ans. (i)



For alkyl halides containing the same alkyl group, the boiling point increases with an increase in the atomic mass of the halogen atom.


Since the atomic mass of Br is greater than that of Cl, the boiling point of bromomethane is higher than that of chloromethane.

Further, for alkyl halides containing the same alkyl group, the boiling point increases with an increase in the number of halides. Therefore, the boiling point of Dibromomethane is higher than that of chloromethane and bromomethane, but lower than that of bromoform.

Hence, the given set of compounds can be arranged in the order of their increasing boiling points as:

Chloromethane < Bromomethane < Dibromomethane < Bromoform.

(ii)



For alkyl halides containing the same halide, the boiling point increases with an increase in the size of the alkyl group. Thus, the boiling point of 1-chlorobutane is higher than that of isopropyl chloride and 1-chloropropane.


Further, the boiling point decreases with an increase in branching in the chain. Thus, the boiling point of isopropyl alcohol is lower than that of 1-chloropropane.

Hence, the given set of compounds can be arranged in the increasing order of their boiling points as:


Isopropyl chloride < 1-Chloropropane < 1-Chlorobutane

7. Which alkyl halide from the following pairs would you expect to react more rapidly by an S_N2 mechanism? Explain your answer.


(i)




(ii)



(iii)




Ans. (i)




2-bromobutane is a 2° alkylhalide whereas 1-bromobutane is a 1° alkyl halide. The approaching of nucleophile is more hindered in 2-bromobutane than in 1-bromobutane. Therefore, 1-bromobutane reacts more rapidly than 2-bromobutane by an S_N2 mechanism.

(ii)




2-Bromobutane is 2° alkylhalide whereas 2-bromo-2-methylpropane is 3° alkyl halide. Therefore, greater numbers of substituents are present in 3° alkyl halide than in 2° alkyl halide to hinder the approaching nucleophile. Hence, 2-bromobutane reacts more rapidly than 2-bromo-2-methylpropane by an S_N2 mechanism.

(iii)



1 - Bromo - 3 - methylbutane (1°)




1 - Bromo - 2 - methylbutane (1°)


Both the alkyl halides are primary. However, the substituent $-\text{CH}_3$ is at a greater distance to the carbon atom linked to Br in 1-bromo-3-methylbutane than in 1-bromo-2-methylbutane. Therefore, the approaching nucleophile is less hindered in case of the former than in case of the latter. Hence, the former reacts faster than the latter by S_N2 mechanism.

8. In the following pairs of halogen compounds, which compound undergoes faster S_N1 reaction?



(i)



(ii)



Ans. (i)




S_N1 reaction proceeds via the formation of carbocation. The alkyl halide (I) is 3° while (II) is 2° . Therefore, (I) forms 3° carbocation while (II) forms 2° carbocation. Greater the stability of the carbocation, faster is the rate of S_N1 reaction. Since 3° carbocation is more stable than 2° carbocation. (I), i.e. 2-chloro-2-methylpropane, undergoes faster S_N1 reaction than (II) i.e., 3-chloropentane.

(ii)




The alkyl halide (I) is 2° while (II) is 1° . 2° carbocation is more stable than 1° carbocation. Therefore, (I), 2-chloroheptane, undergoes faster S_N1 reaction than (II), 1-chlorohexane.


9. Identify A, B, C, D, E, R and R^1 in the following:





Ans.



Since D of D_2O gets attached to the carbon atom to which MgBr is attached, C is




Therefore, the compound R - Br is




When an alkyl halide is treated with Na in the presence of ether, a hydrocarbon containing double the number of carbon atoms as present in the original halide is obtained as product.


This is known as Wurtz reaction. Therefore, the halide, R¹-X, is




Therefore, compound D is



And, compound E is








2 - Methylpropane




Chapter End Question

1. Name the following halides according to IUPAC system and classify them as alkyl, allyl, benzyl (primary, secondary, tertiary), vinyl or aryl halides:






Ans. (i)



2-Chloro-3-methylbutane

(Secondary alkyl halide)


(ii)



3-Chloro-4-methylhexane

(Secondary alkyl halide)


(iii)



1-Iodo-2, 2 -dimethylbutane

(Primary alkyl halide)


(iv)



1-Bromo-3,3-dimethyl-1-phenylbutane

(benzylic halide)


(v)



2-Bromo-3-methylbutane

(Secondary alkyl halide)


(vi)



1-Bromo-2-ethyl-2-methylbutane

(Primary alkyl halide)


(vii)



3-Chloro-3-methylpentane

(Tertiary alkyl halide)


(viii)



3-Chloro-5-methylhex-2-ene

(Vinylic halide)


(ix)



4-Bromo-4-methylpent-2-ene

(Tertiary and Allylic halide)


(x)



1-Chloro-4-(2-methylpropyl) benzene

(Aryl halide)


(xi)



1-Chloromethyl-3-(2, 2-dimethylpropyl) benzene

(benzylic halide)

(xii)



1-Bromo-2-(1-methylpropyl) benzene

(Aryl halide)

2. Give the IUPAC names of the following compounds:

(i) $\text{CH}_3\text{CH}(\text{Cl})\text{CH}(\text{Br})\text{CH}_3$

(ii) $\text{CHF}_2\text{CBrClF}$


(iii) $\text{ClCH}_2\text{C}=\text{CCH}_2\text{Br}$

(iv) $(\text{CCl}_3)_3\text{CCl}$

(v) $\text{CH}_3\text{C}(\text{p-ClC}_6\text{H}_4)_2\text{CH}(\text{Br})\text{CH}_3$


(vi) $(\text{CH}_3)_3\text{CCH}=\text{CClC}_6\text{H}_4\text{I-p}$

Ans. (i)




2-Bromo-3-chlorobutane

(ii)




1-Bromo-1-chloro-1, 2, 2-trifluoroethane

(iii)




1-Bromo-4-chlorobut-2-yne

(iv)




1,1,1,2,3,3,3-heptachloro-2-(trichloromethyl)propane [tris-(trichloromethyl)chloromethane]

(v)



2-Bromo-3, 3-bis(4-chlorophenyl) butane

(vi)



1-chloro-1-(4-iodophenyl)-3, 3-dimethylbut-1-ene

3. Write the structures of the following organic halogen compounds.

(i) 2-Chloro-3-methylpentane

(ii) *p*-Bromochlorobenzene

(iii) 1-Chloro-4-ethylcyclohexane

(iv) 2-(2-Chlorophenyl)-1-iodooctane


(v) Perfluorobenzene

(vi) 4-tert-Butyl-3-iodoheptane

(vii) 1-Bromo-4-sec-butyl-2-methylbenzene


(viii) 1,4-Dibromobut-2-ene

Ans. (i)




2-Chloro-3-methylpentane

(ii)




p-Bromochlorobenzene

(iii)



1-Chloro-4-ethylcyclohexane

(iv)




2-(2-Chlorophenyl)-1-iodooctane

(v)




Perfluorobenzene

(vi)




4-Tert-Butyl-3-iodoheptane

(vii)



1-Bromo-4-sec-butyl-2-methylbenzene

(viii)



1,4-Dibromobut-2-ene

4. Which one of the following has the highest dipole moment?

(i) CH_2Cl_2

(ii) CHCl_3

(iii) CCl_4


Ans. (i)



Dichloromethane (CH_2Cl_2) has highest dipole moment.

$$\mu = 1.60\text{D}$$


(ii)



Chloroform (CHCl_3) has 3C-Cl bonds which opposw resultant 2 dipoles.

$$\mu = 1.03\text{D}$$

(iii)



Carbon tetrachloride (CCl_4) has regular tetrahedral structure so individual dipole gets cancelled.


$$\mu = 0\text{D}$$

CCl_4 is a symmetrical molecule. Therefore, the dipole moments of all four C-Cl bonds cancel each other. Hence, its resultant dipole moment is zero.

As shown in the above figure, in CHCl_3 , the resultant of dipole moments of two C-Cl bonds is opposed by the resultant of dipole moments of one C-H bond and one C-Cl bond. Since the resultant of one C-H bond and one C-Cl bond dipole moments is smaller than two C-Cl bonds, the opposition is to a small extent. As a result, CHCl_3 has a small dipole moment of 1.03 D.

On the other hand, in case of CH_2Cl_2 , the resultant of the dipole moments of two C-Cl bonds is strengthened by the resultant of the dipole moments of two C-H bonds. As a result, CH_2Cl_2 has a higher dipole moment of 1.60 D than CHCl_3 i.e., CH_2Cl_2 has the highest dipole moment.

Hence, the given compounds can be arranged in the increasing order of their dipole moments as:



5. A hydrocarbon C_5H_{10} does not react with chlorine in dark but gives a single monochloro compound $\text{C}_5\text{H}_9\text{Cl}$ in bright sunlight. Identify the hydrocarbon.

Ans. A hydrocarbon with the molecular formula, C_5H_{10} belongs to the group with a general molecular formula C_nH_{2n} . Therefore, it may either be an alkene or a cycloalkane.


Since hydrocarbon does not react with chlorine in the dark, it cannot be an alkene. Thus, it should be a cycloalkane.

Further, the hydrocarbon gives a single monochloro compound, $\text{C}_5\text{H}_9\text{Cl}$ by reacting with chlorine in bright sunlight. Since a single monochloro compound is formed, the hydrocarbon must contain all equivalent H- atoms. Also, as all H-atoms of a cycloalkane are equivalent, the hydrocarbon must be a cycloalkane. Hence, the said compound is cyclopentane.



Cyclopentane (C_5H_{10})


The reactions involved in the question are:



6. Write the isomers of the compound having formula C₄H₉Br.


Ans. There are four isomers of the compound having the formula C₄H₉Br. These isomers are given below.

(a)




1-Bromobutane

(b)




2-Bromobutane

(c)



1-Bromo-2-methylpropane

(d)



2-Bromo-2-methylpropane


7. Write the equations for the preparation of 1-iodobutane from

(i) 1-butanol


(ii) 1-chlorobutane

(iii) but-1-ene.


Ans. (i)



(ii)




(iii)




8. What are ambidentate nucleophiles? Explain with an example.

Ans. Ambidentate nucleophiles are nucleophiles having two nucleophilic sites. Thus, ambidentate nucleophiles have two sites through which they can attack and form bonds.

For example, Cyanide ion, nitrite ion are ambidentate nucleophile.



Nitrite ion can attack through oxygen resulting in the formation of alkyl nitrites. Also, it can attack through nitrogen resulting in the formation of nitroalkanes.




9. Which compound in each of the following pairs will react faster in $\text{S}_{\text{N}}2$ reaction with OH^- ?

(i) CH_3Br or CH_3I


(ii) $(\text{CH}_3)_3\text{CCl}$ or CH_3Cl

Ans. (i) In the $\text{S}_{\text{N}}2$ mechanism, the reactivity of halides for the same alkyl group increases in the order. This happens because as the size increases, the halide ion becomes a better leaving group.



As, CH_3I will react faster because it has low bond dissociation energy and iodine is a better leaving group than CH_3Br in $\text{S}_{\text{N}}2$ reactions with OH^-

(ii)



The S_N2 mechanism is very fast with primary halides and it involves the attack of the nucleophile at the atom bearing the leaving group. But, in case of $(\text{CH}_3)_3\text{CCl}$, the attack of the nucleophile at the carbon atom is hindered because of the presence of bulky substituents on that carbon atom bearing the leaving group. On the other hand, there are no bulky substituents on the carbon atom bearing the leaving group in CH_3Cl . Hence, CH_3Cl reacts faster than $(\text{CH}_3)_3\text{CCl}$ in S_N2 reaction with OH^- .


10. Predict all the alkenes that would be formed by dehydrohalogenation of the following halides with sodium ethoxide in ethanol and identify the major alkene:

(i) 1-Bromo-1-methylcyclohexane


(ii) 2-Chloro-2-methylbutane

(iii) 2,2,3-Trimethyl-3-bromopentane.


Ans. (i)




In the given compound, there are two types of β -hydrogen atoms are present. Thus, dehydrohalogenation of this compound gives only one alkenes.



(ii)




In the given compound, there are two different sets of equivalent α -hydrogen atoms labelled as *a* and *b*. Thus, dehydrohalogenation of the compound yields two alkenes.



Saytzeff's rule implies that in dehydrohalogenation reactions, the alkene having a greater number of alkyl groups attached to a doubly bonded carbon atoms is preferably produced.


Therefore, alkene (I) i.e., 2-methylbut-2-ene is the major product in this reaction.

(iii)



2,2,3-Trimethyl-3-bromopentane

In the given compound, there are two different sets of equivalent β -hydrogen atoms labelled *a* and *b*. Thus, dehydrohalogenation of the compound yields two alkenes.



According to Saytzeff's rule, in dehydrohalogenation reactions, the alkene having a greater number of alkyl groups attached to the doubly bonded carbon atom is preferably formed.

Hence, alkene (I) i.e., 3,4,4-trimethylpent-2-ene is the major product in this reaction.



11. How will you bring about the following conversions?

- (i) Ethanol to but-1-yne
- (ii) Ethane to bromoethene
- (iii) Propene to 1-nitropropane
- (iv) Toluene to benzyl alcohol
- (v) Propene to propyne
- (vi) Ethanol to ethyl fluoride
- (vii) Bromomethane to propanone
- (viii) But-1-ene to but-2-ene





(ix) 1-Chlorobutane to n-octane

(x) Benzene to biphenyl.


Ans. (i)




(ii)




(iii)




(iv)




(v)




(vi)



(vii)




(viii)



(ix)



(x)




Chapter End Question

12. Explain why

- (i) the dipole moment of chlorobenzene is lower than that of cyclohexyl chloride?
- (ii) alkyl halides, though polar, are immiscible with water?

(iii) Grignard reagents should be prepared under anhydrous conditions?

Ans. (i)




In chlorobenzene, the Cl-atom is linked to a sp^2 hybridized carbon atom and stabilized by resonance. In cyclohexyl chloride, the Cl-atom is linked to a sp^3 hybridized carbon atom. Now, sp^2 hybridized carbon has more +-character on three resonating structures therefore it has lower dipole moment than sp^3 hybridized carbon atom. Therefore, the former is more electronegative than the latter. However, there is no such positive charge on cyclohexyl chloride.

Moreover, the -R effect of the benzene ring of chlorobenzene decreases the electron density of the C - Cl bond near the Cl-atom. As a result, the polarity of the C - Cl bond in chlorobenzene decreases. Hence, the dipole moment of chlorobenzene is lower than that of cyclohexyl chloride.

(ii) To be miscible with water, the solute-water force of attraction must be stronger than the solute-solute and water-water forces of attraction. Alkyl halides are polar molecules and so held together by dipole-dipole interactions but can not form H-bonds with water neither it can break H-bonds between water molecules. Similarly, strong H-bonds exist between the water molecules. The new force of attraction between the alkyl halides and water molecules is weaker than the alkyl halide-alkyl halide and water-water forces of attraction. Hence, alkyl halides (though polar) are immiscible with water.

(iii) Grignard reagents are very reactive. In the presence of moisture, they react to give alkanes.



Therefore, Grignard reagents should be prepared under anhydrous conditions.

13. Give the uses of freon 12, DDT, carbon tetrachloride and iodoform.

Ans. Uses of Freon - 12

Freon-12 (dichlorodifluoromethane, CF_2Cl_2) is commonly known as CFC. It is used as a refrigerant in refrigerators and air conditioners. It is also used in aerosol spray propellants such as body sprays, hair sprays, etc. However, it damages the ozone layer. Hence, its manufacture was banned in the United States and many other countries in 1994.

Uses of DDT

DDT (*p, p*-dichlorodiphenyltrichloroethane) is one of the best known insecticides. It is very effective against mosquitoes and lice. But due to its harmful effects, it was banned in the United States in 1973.

Uses of carbontetrachloride (CCl_4)









- (i) It is used for manufacturing refrigerants and propellants for aerosol cans.
- (ii) It is used as feedstock in the synthesis of chlorofluorocarbons and other chemicals.
- (iii) It is used as a solvent in the manufacture of pharmaceutical products.
- (iv) Until the mid 1960's, carbon tetrachloride was widely used as a cleaning fluid, a degreasing agent in industries, a spot reamer in homes, and a fire extinguisher because vapours are heavy cut off supply of air.

Uses of iodoform (CHI_3)



Iodoform was used earlier as an antiseptic, but now it has been replaced by other formulations-containing iodine-due to its objectionable smell. The antiseptic property of




iodoform is only due to the liberation of free iodine when it comes in contact with the skin.

14. Write the structure of the major organic product in each of the following reactions:



Ans.





(viii)




15. Write the mechanism of the following reaction:



Ans. The given reaction is a nucleophilic substitution reaction.



The given reaction is an $\text{S}_{\text{N}}2$ reaction. In this reaction, CN^- acts as the stronger nucleophile and attacks the carbon atom to which Br which is a weaker nucleophile is attached. CN^- ion is an ambident nucleophile and can attack through both C and N. In this case, it attacks through the C-atom.




16. Arrange the compounds of each set in order of reactivity towards S_N2 displacement:

(i) 2-Bromo-2-methylbutane, 1-Bromopentane, 2-Bromopentane

(ii) 1-Bromo-3-methylbutane, 2-Bromo-2-methylbutane, 3-Bromo-2-methylbutane

(iii) 1-Bromobutane, 1-Bromo-2,2-dimethylpropane, 1-Bromo-2-methylbutane, 1-Bromo-3-methylbutane.

Ans. (i)



An S_N2 reaction involves the approaching of the nucleophile to the carbon atom to which the leaving group is attached. When the nucleophile is sterically hindered, then the reactivity towards S_N2 displacement decreases. Due to the presence of substituents, hindrance to the approaching nucleophile increases in the following order.

1-Bromopentane < 2-bromopentane < 2-Bromo-2-methylbutane


Hence, the increasing order of reactivity towards S_N2 displacement is:

2-Bromo-2-methylbutane < 2-Bromopentane < 1-Bromopentane


(ii)



1-Bromo-3-methylbutane
(1°)



2-Bromo-2-methylbutane
(3°)



2-Bromo-3-methylbutane
(2°)

Since steric hinderance in alkyl halides increases in the order of $1^\circ < 2^\circ < 3^\circ$, the increasing order of reactivity towards S_N2 displacement is given as-





$$3^\circ < 2^\circ < 1^\circ.$$

Hence, the given set of compounds can be arranged in the increasing order of their reactivity towards S_N2 displacement as:




[2-Bromo-3-methylbutane is incorrectly given in NCERT]

(iii)



The steric hinderance to the nucleophile in the S_N2 mechanism increases with a decrease in the distance of the substituents from the atom containing the leaving group. Further, the steric hinderance increases with an increase in the number of substituents. Therefore, the increasing order of steric hindrances in the given compounds is as below:





Hence, the increasing order of reactivity of the given compounds towards S_N2 displacement

is:

1-Bromo-2, 2-dimethylpropane < 1-Bromo-2-methylbutane < 1-Bromo-3- methylbutane < 1-Bromobutane

17. Out of $C_6H_5CH_2Cl$ and $C_6H_5CHClC_6H_5$, which is more easily hydrolysed by aqueous KOH?


Ans.




Hydrolysis by aqueous KOH proceeds through the formation of carbocation. If carbocation is stable, then the compound is easily hydrolyzed by aqueous KOH. Now $C_6H_5CH_2Cl$, forms 1° -carbocation, while $C_6H_5CHClC_6H_5$ forms 2° -carbocation, which is more stable than 1° -carbocation. Hence $C_6H_5CHClC_6H_5$, is hydrolyzed more easily than $C_6H_5CH_2Cl$ by aqueous KOH.

18. *p*-Dichlorobenzene has higher m.p. and lower solubility than those of *o*- and *m*-isomers. Discuss.


Ans.



p-Dichlorobenzene



o-Dichlorobenzene



m-Dichlorobenzene

p-Dichlorobenzene is more symmetrical than *o*- and *m*-isomers. For this reason, it fits more

closely and easily in the crystal lattice than *o*- and *m*-isomers. Therefore, more energy is required to break the crystal lattice of *p*-dichlorobenzene. As a result, *p*-dichlorobenzene is symmetrical and has a higher melting point and lower solubility than *o*- and *m*-isomers due to strong force of attraction in crystal.

19. How the following conversions can be carried out?


- (i) Propene to propan-1-ol**
- (ii) Ethanol to but-1-yne**
- (iii) 1-Bromopropane to 2-bromopropane**
- (iv) Toluene to benzyl alcohol**
- (v) Benzene to 4-bromonitrobenzene**
- (vi) Benzyl alcohol to 2-phenylethanoic acid**
- (vii) Ethanol to propanenitrile**
- (viii) Aniline to chlorobenzene**
- (ix) 2-Chlorobutane to 3, 4-dimethylhexane**
- (x) 2-Methyl-1-propene to 2-chloro-2-methylpropane**
- (xi) Ethyl chloride to propanoic acid**
- (xii) But-1-ene to n-butyliodide**
- (xiii) 2-Chloropropane to 1-propanol**
- (xiv) Isopropyl alcohol to iodoform**
- (xv) Chlorobenzene to *p*-nitrophenol**
- (xvi) 2-Bromopropane to 1-bromopropane**
- (xvii) Chloroethane to butane**

(xviii) Benzene to diphenyl


(xix) *tert*-Butyl bromide to isobutyl bromide

(xx) Aniline to phenylisocyanide


Ans. (i)




(ii)



(iii)




(iv)




(v)




(vi)




(vii)



(viii)




(ix)




(x)




(xi)




(xii)




(xiii)




(xiv)




(xv)







(xvi)




(xvii)




(xviii)



(xix)




(xx)




20. The treatment of alkyl chlorides with aqueous KOH leads to the formation of alcohols but in the presence of alcoholic KOH, alkenes are major products. Explain.

Ans. In an aqueous solution, KOH almost completely ionizes to give OH⁻ ions. OH⁻ ion is a strong nucleophile, which leads the alkyl chloride to undergo a nucleophilic substitution reaction to form alcohol.



On the other hand, an alcoholic solution of KOH contains alkoxide (RO⁻) ion, which is a strong base. Thus, it can abstract a hydrogen from the β-carbon of the alkyl chloride and form an alkene by eliminating a molecule of HCl.




OH⁻ ion is a much weaker base than RO⁻ ion. Also, OH⁻ ion is highly solvated in an aqueous solution and as a result, the basic character of OH⁻ ion decreases. Therefore, it cannot abstract a hydrogen from the β-carbon.

Concept Insight: OH⁻ is a strong nucleophile but weaker base than RO⁻

21. Primary alkyl halide C₄H₉Br (a) reacted with alcoholic KOH to give compound (b). Compound (b) is reacted with HBr to give (c) which is an isomer of (a). When (a) is reacted with sodium metal it gives compound (d), C₈H₁₈ which is different from the compound formed when n-butyl bromide is reacted with sodium. Give the structural



formula of (a) and write the equations for all the reactions.

Ans. There are two primary alkyl halides having the formula, C_4H_9Br . They are *n*-butyl bromide and isobutyl bromide.




Therefore, compound (a) is either *n*-butyl bromide or isobutyl bromide.

Now, compound (a) reacts with Na metal to give compound (b) of molecular formula, C_8H_{18} , which is different from the compound formed when *n*-butyl bromide reacts with Na metal. Hence, compound (a) must be isobutyl bromide.




Thus, compound (d) is 2, 5-dimethylhexane.

It is given that compound (a) reacts with alcoholic KOH to give compound (b). Hence, compound (b) is 2-methylpropene.




Also, compound (b) reacts with HBr to give compound (c) which is an isomer of (a). Hence, compound (c) is 2-bromo-2-methylpropane.




22. What happens when

- (i) n-butyl chloride is treated with alcoholic KOH,
- (ii) bromobenzene is treated with Mg in the presence of dry ether,
- (iii) chlorobenzene is subjected to hydrolysis,
- (iv) ethyl chloride is treated with aqueous KOH,
- (v) methyl bromide is treated with sodium in the presence of dry ether,
- (vi) methyl chloride is treated with KCN.

Ans. (i) When n - butyl chloride is treated with alcoholic KOH, the formation of but - 1 - ene takes place. This reaction is a dehydrohalogenation reaction.




(ii) When bromobenzene is treated with Mg in the presence of dry ether, phenylmagnesium bromide is formed.




(iii) Chlorobenzene does not undergo hydrolysis under normal conditions. However, it


undergoes hydrolysis when heated in an aqueous sodium hydroxide solution at a temperature of 623 K and a pressure of 300 atm to form phenol.



(iv) When ethyl chloride is treated with aqueous KOH, it undergoes hydrolysis to form ethanol.



(v) When methyl bromide is treated with sodium in the presence of dry ether, ethane is formed. This reaction is known as the Wurtz reaction.



(vi) When methyl chloride is treated with KCN, it undergoes a substitution reaction to give methyl cyanide.

